

Magnetic resonance imaging of hip joint cartilage and labrum

Christoph Zilkens,¹ Falk Miese,² Marcus Jäger,³ Bernd Bittersohl,¹ Rüdiger Krauspe¹

¹ Department of Orthopaedic Surgery, University Hospital of Düsseldorf, Germany

² Department of Radiology, University Hospital of Düsseldorf, Germany

³ Department of Orthopaedic Surgery, University Hospital of Essen, Germany

Abstract

Hip joint instability and impingement are the most common biomechanical risk factors that put the hip joint at risk to develop premature osteoarthritis. Several surgical procedures like periacetabular osteotomy for hip dysplasia or hip arthroscopy or safe surgical hip dislocation for femoroacetabular impingement aim at restoring the hip anatomy. However, the success of joint preserving surgical procedures is limited by the amount of pre-existing cartilage damage. Biochemically sensitive MRI techniques like delayed Gadolinium Enhanced MRI of Cartilage (dGEMRIC) might help to monitor the effect of surgical or non-surgical procedures in the effort to halt or even reverse joint damage.

Introduction

Osteoarthritis (OA) of the hip joint is a major cause for disability and pain in the adult population of developed countries.¹ Instability and impingement or combinations of instability and impingement are the most important mechanical factors that put the hip joint at risk of developing early OA.² Childhood diseases like hip dysplasia (instability), Legg-Calve-Perthes disease (static impingement) or slipped capital femoral epiphysis (dynamic impingement) are major etiologic contributors to the development of early hip OA.³ While 50 years ago it was assumed that 50% of hip OA was not attributable to anatomic deformities (idiopathic OA),⁴ nowadays some authors suspect that more than 90% of hip OA is due to instability or impingement.²

In order to diagnose and treat patients with FAI or hip dysplasia according to the disease severity, adequate knowledge of magnetic resonance imaging of the hip joint pathology is mandatory.

Hip joint anatomy

The hip joint is large, has to bear a lot of weight and its stability is provided by its rigid ball-and-socket or nut-configuration as well as the surrounding strong ligaments and muscles. The acetabular cartilage is horse-shoe-shaped with a central part without cartilage coverage that does not articulate with the femoral head (fossa acetabuli). Within the fossa, fatty tissue and the ligamentum teres are imaged on MRI. The femoral head is completely covered with hyaline cartilage except for the insertion of the ligamentum teres. The hip joint cartilage is thin in comparison to other joints with the maximum thickness ventrocranially at the acetabulum and ventrolaterally on the femoral head.

The joint capsule is strengthened by 3 ligaments: the iliofemoral ligament is the strongest ligament of the 3 and originates from between the anterior inferior iliac spine and the acetabular rim and inserts along the anterior portion of the intertrochanteric line and greater trochanter. It assists in the maintenance of an erect posture without much muscular activity. The pubofemoral ligament originates from the ramus superior ossis pubis and inserts anterolaterally in the joint capsule while the ischiofemoral ligament is dorsally, originating from the ischium and going horizontally inserting on the upper limit of the intertrochanteric line.⁵

With increasing interest in hip arthroscopy, the role of the ligamentum teres as a secondary contributor to hip stability⁶ is under re-investigation: lesions of the ligamentum teres have gained attentiveness through hip arthroscopy and have been described in up to 15% of hip arthroscopy patients and as a common cause of hip pain in athletes.⁶⁻⁹ The ligamentum teres (Figure 11H) arises from the transverse acetabular ligament and is attached to the periosteum by to fascicles along the ischial and pubic margins of the acetabular notch.⁶ The acetabular labrum is a sealing rim around the hip joint that consists of fibrocartilaginous collagen fibers attached to the acetabulum and contiguous with the transverse acetabular ligament. The functions of the labrum comprise an increase of the acetabular volume, dissipation of force across the hip, facilitation of synovial lubrication, compensation for minor joint incongruities and dissipation of contact forces encountered by the hip joint.¹⁰⁻¹⁵ The capsular side of the labrum consists of dense connective tissue, whereas the articular side is composed of fibrocartilage.¹⁶ Without intrinsic vasculature the blood supply is provided by the capsule and synovium.¹⁶⁻¹⁹ Its nociceptive and proprioceptive function are still under investigation. Different types of corpuscles represent pressure receptors, receptors of deep sensation and temperature sensation while free nerve endings are pain recep-

Correspondence: Christoph Zilkens, Department of Orthopaedic Surgery, University Hospital Düsseldorf, Moorenstrasse 5, 40225 Düsseldorf, Germany. Tel. 0211.8118314 - Fax: 0211.8117965. E-mail: christoph.zilkens@med.uni-duesseldorf.de

Received for publication: 27 May 2011.
Accepted for publication: 10 June 2011.

This work is licensed under a Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0).

©Copyright C. Zilkens et al., 2011
Licensee PAGEPress, Italy
Orthopedic Reviews 2011; 3:e9
doi:10.4081/or.2011.e9

tors.^{10,17,20} Other than the nociceptive and proprioceptive impairment, a torn labrum will cause a reduction in the described mechanical sealing support thus not anymore maintaining the synovial fluid for force distribution, smooth gliding surface, and nutrition and thus resulting in more cartilage damage.²¹

Rule out other factors of hip pain

In contrast to other joints and due to its anatomic position the hip joint is not always easy to examine and pain around the hip joint might be due to other factors than labral or cartilage damage due to FAI or dysplasia.

Avulsion fractures, insufficiency fractures, osteoporotic or pathologic fractures and tumors around the hip joint have to be ruled out as cause for hip pain. Chronic inflammatory arthritis including rheumatoid arthritis might be accompanied with morning stiffness and other systemic manifestations of the disease. Lumbar radiculopathy and lumbar spinal stenosis might mimic hip pain. Intrapment of the lateral femoral cutaneous nerve might cause meralgia paresthetica with pain or numbness on the lateral aspect of the hip and thigh. Lose intraarticular bodies, gout or pseudogout, synovitis or acute bacterial arthritis have to be ruled out as reason for hip pain. Piriformis syndrome is referred to an irritation of the sciatic nerve by the piriformis muscle. Iliotibial band syndrome might radiate along the lateral thigh and cause an external snapping hip, in contrast to the internal snapping hip that is caused by the iliopsoas muscle. Sports hernia or athletic pubalgia are occult hernias caused by weakness or tear of the posterior inguinal wall without recognizable hernia, Gilmore's groin with tear in the external oblique aponeurosis, conjoined tendon and dehiscence between the conjoined tendon and the inguinal ligament as well as injury at the insertion of the rectus abdominis muscle, avulsion of the internal oblique muscle or tearing within the internal or external oblique aponeurosis or muscle (Figure 1).²²⁻²⁴

Hip abductors have been compared to the rotator cuff of the shoulder^{25,26} and gluteus medius tendinous or muscular pathologies including fatty degeneration may be graded and treated according to rotator cuff pathologies.²⁷

Other reasons for referred hip pain might be synovitis or mechanical blockade or the sacroiliac joint, osteitis pubis, muscle injuries and enthesiopathies of the adductors, iliopsoas or hamstrings. Chronic microtrauma and injury to the adductors might be caused by an externally rotating *cam-avoidance* gait pattern in cam-FAI-patients.^{28,29} A *sports hip triad* has been described recently, consisting of a labral tear, adductor strain and rectus strain.³⁰

Bursae might be inflamed and swollen and might be mistaken for tumors or cysts. Bursae commonly affected by acute or chronic bursitis are the greater trochanteric bursa, the iliopsoas bursa (= iliopsoas bursa) and the ischiogluteal bursa (Figure 2).

The iliopsoas bursa is the biggest bursa around the hip joint and might communicate with the hip joint in 15% of the people. That is why in MR-artrograms contrast agent might extend into the iliopsoas (Figure 3).

Therapy of avascular necrosis (AVN) of the femoral head depends on the stage of the disease. MRI has reported sensitivities and specificities as high as 100% for the detection of ON (Figure 4).³¹ Treatment strategies for AVN depend on the stage of the disease that might be classified by the Association Research Circulation Osseous (ARCO).^{32,34} Since joint preserving procedures for advanced stages of AVN are limited, early diagnosis and effective treatment are necessary.^{35,36} The vasoactive, stable prostacyclin analogue iloprost is approved for therapy of critical limb ischemia due to peripheral arteriosclerotic obliterative disease and diabetic angiopathy as well as an inhalative for patients with pulmonary arterial hypertension.³⁷ Our group and others use iloprost for the treatment of early stages of AVN.³⁸⁻⁴¹

Labrum

The healthy labrum has a triangular shape with sharp margins and continuous attachment to the acetabular rim and cartilage (chondrolabral junction).⁴² The labrum is contiguous with the transverse acetabular ligament, which appears cuboid and marks the medial-inferior part of fossa acetabuli. A labral tear shows increased intra-substance signal with in labral detachment from the acetabular rim, synovial-fluid-intensity signal will undermine the labrum. Labral tears are typically located antero-superiorly. A degenerated labrum appears clumsy with intralabral signal alteration due to mucoid degeneration (Figure 5).

In order to achieve useful images, high MR resolution and contrast to noise ratio are

required. Non-contrast MRI is used for the evaluation of bone, necroses, tumors, muscles and marrow space. It seems to be unreliable for detecting more subtle lesions. Mintz et al. found a sensitivity of 96%, a specificity of 33% and an overall accuracy of 94% for the detection of labral tears at 1.5T.⁴³ Sundberg et al. found comparable results for the detection of labral tears comparing 3-T non-artrographic with 1.5-T artrographic techniques.⁴⁴ With the studies available today, non-contrast MRI is not optimal in the evaluation of cartilage and labrum. In the future and with more sophisticated hardware and software as well as the availability of higher field strength machines, this may change. Direct Magnetic resonance arthrography (d-MRA) after the intra-articular injection of gadolinium-based contrast agent has emerged as the standard method for the evaluation of labrum and cartilage.⁴⁵⁻⁴⁸ Approximately 10-20 mL of contrast agent is injected into the hip joint under fluoroscopic guidance, followed by MRI within approxi-




Figure 1. Common locations of tendinosis and sprain in the athlete.




Figure 2. Bursae of clinical relevance.




Figure 3. Bursitis and distension of a bursa iliopsoas.




Figure 4. AVN ARCO I.




Figure 5. Torn Labrum, suture during surgical hip dislocation.




Figure 6. Intra articular contrast administration.

mately 30 minutes (Figure 6).⁴⁹ The intraarticular contrast agent increases the spatial resolution and causes a capsular distension with separation of capsule, labrum and osteochondral structures. The contrast agent can fill into labral and cartilage clefts. Compared to hip arthroscopy as gold standard, d-MRA is reported to have sensitivities of 63-100%, specificities of 44-100% and accuracy values of 65-96%.^{50,55} For the detection of labral tears, the inter-observer reliability has been reported to be moderate.^{50,56} With 2 dimensional in contrast to 3 dimensional MRI techniques, the assessment of thickness and orientation of the acetabular lesion was not optimal.^{45,50} For indirect MRA (i-MRA) of the hip joint, gadolinium-containing contrast agent is administered intravenously, followed by a delay with or without physical activity. The contrast agent will distribute into the joint, enhancing synovial fluid and providing greater contrast as well as distension of the capsule, allowing for the interpretation of labrum and cartilage.^{49,57,61} In one study comparing i-MRA and d-MRA, i-MRA showed a sensitivity of 88% and an accuracy of 90%.⁶² Byrd *et al.* demonstrated that D-MRA was much more sensitive in the detection of various lesions, however, arthroscopy demonstrated that d-MRA was interpreted falsely positive twice as much compared to i-MRA (Figure 7).⁶³ One major advantage for the d-MRA is the possibility to perform a diagnostic infiltration of the hip joint at the same time as contrast agent administration: it has been shown that the reduction of pain after intra-articular administration of a local anesthetic is a 90% reliable indicator of intraarticular pathology.⁶³ However, the informational value of this probe-infiltration of the hip joint is diminished by the fact that with administration of 20 mL of contrast agent the joint capsule is distended, causing pain itself. Advantages of i-MRA versus d-MRA comprise: the lesser risk of vascular or nerve injury by the injection; the absence of radiation through fluoroscopy; the reduced resource and time intensity as well as reduced logistical effort.

Cartilage

Hip joint cartilage is thin and bony hip anatomy is complex with the shape of the head being more or less spherical. Cartilage lesion assessment is not as well established as labrum lesion assessment. Non-contrast techniques to describe cartilage changes revealed low diagnostic efficiency with sensitivities of less than 50%.⁶⁴ Mintz *et al.*⁴³ described a low reliability in classifying cartilage according to cartilage thickness and signal intensity changes according to the Outerbridge Score.⁶⁵ In a study of Schmid *et al.*, the sensitivity of cartilage grading was only 47%.⁵⁶ Overall the cartilage diagnosis in the hip joint is limited so far and no reliable staging and grading system has been

established.^{50,56,66,67} The articular cartilage can be graded with a modification of the classification system of Outerbridge (Table 1).⁴³

Femoroacetabular Impingement

The concept of femoroacetabular impingement (FAI) as a major contributor to the development of premature hip OA has been recognized and accepted all over the world. Table 2 demonstrates the remarkable number of publications in PUBMED concerning *femoroacetabular impingement* within the past decade. The cam-lesion is the reduced head-neck offset and bashes against labrum and acetabular cartilage during flexion and internal rotation. This mechanism may cause cartilage delamination from the subchondral bone and labrum. This carpet phenomenon is located mostly in the anterosuperior region of the acetabulum.⁶⁸⁻⁷⁰ as well as causing intraarticular cartilage damage. In pincer FAI, the acetabulum might be too deep globally or locally, causing an abutment of the femoral neck against the acetabulum so that the labrum might be damaged prior to cartilage damage.⁷¹⁻⁷⁵ Further causes for FAI are rotational anomalies with reduced femoral neck anteversion and/or reduced acetabular retroversion^{72,76} or a focal overcoverage after periacetabular osteotomy (PAO) (*Bernese Disease*).⁷⁷ In many cases patients show pincer and cam deformities (Figure 8). Untreated FAI can lead to premature osteoarthritis (OA)^{69,83} and surgical intervention by open surgical dislocation of the hip, arthroscopy or combined approaches may be warranted. Surgical treatment is associated with positive medium- and long-term outcome. A comparison of the three therapy methods is difficult due to the different outcome measures employed. Studies directly comparing the approaches are warranted to distinguish more clearly between the different treatment options.⁷⁸ As in surgery for hip dysplasia, the outcome of surgery depends on the quantity of pre-existing OA with poor results in patients with advanced degenerative changes. Beck *et al.* described after favourable results after open or arthroscopic FAI-surgery in particular in the subgroup of patients without advanced OA.^{69,83} Therefore in FAI-as well as in hip dysplasia patients it is of great importance to identify early stages of cartilage degeneration to be able to identify patients that will profit from osteo- and/or chondroplastic types of surgery.

Diagnosis of FAI

Diagnosis of FAI is based on clinical findings, standard x-rays (anteroposterior and lateral) and MRI. Plain radiographs are often inadequate in underrepresenting the extent of head-neck pathology.⁷⁹ Due to the importance of detecting the extent of the deformity as well as early cartilage and labral lesions, MRI is the standard tool for diagnosis of FAI.⁴⁶ Further-




Figure 7. Comparison of i-MRA and d-MRA in one Patient with FAI, cartilage damage and torn labrum.





Figure 8. Radial image of a patient with mixed FAI demonstrating a deep socket and a bump deformity. Corresponding labral tear and paralabral cyst.

Table 1. Modified Outerbridge classification for cartilage damage.

	Grade	Macroscopy	MRI
	0	Normal cartilage	Normal cartilage
	1	Rough surface; chondral softening	Inhomogenous, high signal; surface intact
	2	Irregular surface defects; <50% of cartilage thickness	Superficial ulceration, fissuring, fibrillation; <50% of cartilage thickness
	3	Loss of >50% of cartilage thickness	Ulceration, fissuring, fibrillation; > 50% of the depth of cartilage
	4	Cartilage loss	Full thickness chondral wear with exposure of subchondral bone

Table 2. Publications concerning FAI.



more, it is becoming clear that standard coronal, axial and sagittal MR views are less reliable than radially reconstructed planes perpendicular to the acetabular labrum in detecting early degenerative pathologies of the hip.^{46,80}

For the assessment of the femoral head-neck morphology, radial reconstructions along the femoral neck axis are described^{54,81,82} that improve the understanding of the FAI pathomechanism and correlate well with the prediction of an FAI and intra-operative findings.⁸³ These imaging techniques are increasingly recognized as an important tool for morphologic assessment of FAI as well as improved techniques to detect early labral and chondral damage in the hip (Figure 9).⁸⁴

Measurements in FAI

On MRI, different parameters defining FAI can be measured: alpha-angle, head-neck-offset, acetabular depth and acetabular version (Figure 10). Easiest to measure and most important is the alpha-angle of Nötzli⁸⁶ that can be measured as described by Pfirman:⁶⁷ the angle is measured between an axis parallel to the femoral neck and passing through the narrowest portion of the femoral neck, and an axis passing through the point where the head contour passes into the metaphysis as shown in Figure 11K. An angle of more than 55 degrees is indicative of cam deformity. An interval of 30° among the radial reformats should be used to assess alpha angle. The acetabular coverage might be measured by assessing the acetabular depth within in axial reformation. The depth is expressed as distance between a line drawn among anterior and posterior acetabular horn and the center of the femoral head. The acetabular version can be measured on axial 2D T1 weighted images through the acetabular roof, when on the image superiorly where anterior- and posterior rim become apparent. However, acetabular version is better estimated on plain ap radiographs.

Hip dysplasia

In contrast to the FAI, the labrum is typically thick and tears or dissociations are often further dorsolaterally. Chronic overloading of the labrum causes mukoid degeneration and cysts. In hip dysplasia, cartilage damage is more globally than in FAI,⁸⁷ although intra-operative findings show, that cartilage damage occurs predominantly in the antero-superior quadrant both in DDH and in FAI.^{69,71,88} Figure 12 shows the hip joint of a 17 year old patient with symptomatic labral tear and hip dysplasia.

MRI appearance of normal and pathologic features

Labral shape can differ from small and sharp to thick and round or even absent. Increased signal within the labrum is found in sympto-




Figure 9. Radial sequences.




Figure 10. Radial images in a hip dysplasia patient.

matic as well as asymptomatic patients. A poor histologic correlation is reported for these MRI-findings.⁸⁹ Figure 11A shows an intralabral cyst in a 24 years old asymptomatic woman. Figure 11B shows a torn labrum in a symptomatic patient that profited from intra-articular lidocain-injection. Obvious perilabral cysts are shown in Figure 11E in a 28 years old woman with extensive hip dysplasia. Sublabral sulcus or recessus (Figure 11F) are reported to be present in about 25% of patients without pathological meaning^{90,91} while other investigators found no evidence of a normal sublabral sulcus.^{48,92} Perilabral recesses (Figure 11G) can mimic cysts or be mistaken for a labral tear. D-MRA helps appreciating the recess in contrast to i-MRA (Figure 7).^{48,93} Figure 11D shows an os ad acetabuli in a symptomatic FAI-patient. The os ad acetabuli is frequently associated with FAI and might be due to a nonunion of secondary acetabular ossification centers, ossifications of the labrum or incomplete healing of rim fractures.⁹⁴ Supra-acetabular fossae (Figure 11C) appear as additional cavity anterosuperiorly and can be mistaken for osteochondral defects or osteochondrosis dissecans.⁹⁵ Lesions of the ligamentum teres (Figure 11H) have gained attentiveness

through hip arthroscopy and have been described in up to 15% of hip arthroscopy patients and as a common cause of hip pain in athletes. Plicae are embryologic remnants in synovial joints that are often symptomatic in the knee joint, whereas in the hip joint, reports are anecdotic. Fu *et al.*⁹⁶ describe 3 locations for plicae: labral, ligamentous and neck plicae. The pectinofoveal fold is a band that runs parallel to the inferior neck (Figure 11I) with an incidence of 95% in MRI and 99% in hip arthroscopy,⁹⁷ this structure should be regarded as normal and distinguished from pathologic and symptomatic plicae. Slipped capital femoral epiphysis (SCFE)^{98,99} might cause cam impingement and early OA. Figure 11J shows the MRI of a 39-year old woman with advanced OA with osteophytes and capital drop in the long term follow up after SCFE. The alpha angle is added to Figure 11K as mentioned above. Herniation pits (Figure 11L) are fibro-cystic changes along the anterior head-neck-junction that are speculated to be second to FAI.^{100,101}

Biochemical Imaging

Even high field MRI machines image fairly late events while minor changes in cartilage




Figure 11. MRI appearance of normal and pathologic features.

degeneration or regeneration cannot be monitored. Biochemical or molecular imaging of cartilage offers the perspective of closely watching into the cartilage structure. Thus, the real amount of cartilage damage can be visualized and the effect of surgical or non-surgical intervention may be observed. Different biochemical imaging methods are able to visualize cartilage quality in measuring collagen or glycosaminoglycan (GAG) content of cartilage. The imaging modalities can be conducted at regular MRI machines, using the contrast agent, that is also employed on a routine basis. Acquisition time is not much higher than standard morphological sequences. However, the post-processing of the images is still fairly sophisticated and time-consuming, preventing these new and promising techniques to become incorporated into clinical routine. GAGs are proteins of the extracellular cartilage matrix that make out more than 90% of the negative cartilage charge. GAG are lost early in the development of OA¹⁰² and might be replenished in cartilage regeneration. Delayed Gadolinium Enhanced MRI of Cartilage (dGEMRIC) takes advantage of this fact: after intra-articular or intra-venous injection, the negatively charged contrast agent gadolinium-diethylene triamine pentaacetic acid (Gd-DTPA²⁻) penetrates into the cartilage in a reciprocal proportional manner to the content of




Figure 12. Pre-OP x-ray.




Figure 13. Histo and OCT.

GAG within the cartilage. The contrast agent within the cartilage causes a reduction of T1-time that can be measured in MRI. The

dGEMRIC index or $T1_{Gd}$ represents the GAG content within cartilage and high $T1_{Gd}$ values are supposed to be found in healthy cartilage

whereas low $T1_{Gd}$ values are found in degenerated cartilage, due to the higher amount of Gd-DTPA²⁻ within the cartilage. After i.a. or i.v. administration of contrast agent, a delay of 30 to 60 Minutes is warranted before the MRI is performed.¹⁰³ T1 relaxation times that are investigated are: T10 (i.e. T1 prior to contrast administration), $T1_{Gd}$ (post-contrast T1) and $\Delta R1$ that defines the difference in relaxation rate ($R1 = 1/T1$) between $T1_0$ and $T1_{Gd}$ measurements ($1/T1_{Gd} - 1/T1_0$). Some authors found out that $\Delta R1$ is a more precise parameter to reflect the Gd-DTPA²⁻ concentration within cartilage as.¹⁰⁴⁻¹⁰⁶ Bittersohl *et al.* evaluated $T1_{Gd}$ and $\Delta R1$ in two different radiographic grades of hip osteoarthritis in symptomatic FAI patients.¹⁰⁷ Asymptomatic young-adult volunteers served as control. A high correlation between $T1_{Gd}$ and $\Delta R1$ in all study groups could be observed. Based on these results, we conclude that $T1_{Gd}$ assessment is sufficient and a further pre-contrast imaging is not necessary. However, some circumstances require the calculation of $\Delta R1$ for accurate GAG evaluation including follow-up of cartilage repair therapy where T10 values may differ especially in the early postoperative stages post-surgery.^{105,106,108} Several clinical studies have been conducted so far to evaluate hip joint cartilage using dGEMRIC: in his classic report, Kim *et al.* report the diagnostic potential of dGEMRIC for assessment of early OA in patients with hip dysplasia.⁵⁹ Tiderius *et al.* evaluated the time course of T1 values after Gd-DTPA²⁻ injection with hip dysplasia and early signs of OA.¹⁰⁹ The same group investigated 47 patients undergoing a Bernese periacetabular osteotomy (PAO) for the treatment of hip dysplasia.¹¹⁰ Multivariate analysis identified the dGEMRIC index as the most important predictor of failure of the osteotomy. Still the same group retrospectively analyzed 37 symptomatic hips with FAI⁶⁰ and suggested that dGEMRIC may be a useful technique for diagnosis and staging of early osteoarthritis in hips with impingement. Pre-Arthritic deformities after SCFE and Legg-Calve-Perthes disease were evaluated using dGEMRIC.¹¹¹⁻¹¹⁵ GEMRIC may depict the complex damage pattern of hip joint cartilage spatially and qualitatively better than other radiographic methods. The limitation of these studies using 2-D sequences was that only coronal T1 maps could be obtained. However radial evaluation around the hip joint, which is standard in morphologically MRI or MRA, is essential for the detection of cartilage pathologies for.⁴⁶ Recently, fast T1 assessment using dual flip angle (FA) gradient echo (GRE) has been validated and was used in-vivo enabling faster imaging times and three-dimensional (3D) dGEMRIC.^{108,116} This technique utilizes inline T1 measurement and allows for faster imaging. Bittersohl *et al.* proved this technique

to be a reliable instrument in the assessment of asymptomatic hip joint cartilage.¹¹⁷ In a pilot study Bittersohl *et al.* proved the feasibility of cartilage assessment in symptomatic FAI patients using intra-articular delayed Gadolinium Enhanced MRI of Cartilage (ia-dGEMRIC).¹¹⁸ In another study Bittersohl *et al.*¹¹⁹ found that mapping with both iv-dGEMRIC and ia-dGEMRIC demonstrated obvious differences between various grades of cartilage degeneration. In ongoing studies we evaluate sequences histologically: in patients that are scheduled for a total hip endoprosthesis, an in vivo and postoperatively *in vitro* scan of the hip joint cartilage is performed. Both scans can be combined and evaluated histologically. Different sequences are then subject to further immunohistochemical analyses as well as optical coherence tomography. (Figures 12 and 13).

References

1. Dagenais S, Garbedian S, Wai EK. Systematic review of the prevalence of radiographic primary hip osteoarthritis. Clin Orthop Relat Res 2009;467:623-37.
2. Millis MB, Kim YJ. Rationale of osteotomy and related procedures for hip preservation: a review. Clin Orthop Relat Res 2002;108-21.
3. Kim YJ, Ganz R, Murphy SB, et al. Hip joint-preserving surgery: beyond the classic osteotomy. Instr Course Lect 2006;55:145-58.
4. Lloyd-Roberts GC. Osteoarthritis of the hip; a study of the clinical pathology. J Bone Joint Surg Br 1955;37:8-47.
5. Fuss FK, Bacher A. New aspects of the morphology and function of the human hip joint ligaments. Am J Anat 1991;192:1-13.
6. Bardakos NV, Villar RN. The ligamentum teres of the adult hip. J Bone Joint Surg Br 2009;91:8-15.
7. Gray AJ, Villar RN. The ligamentum teres of the hip: an arthroscopic classification of its pathology. Arthroscopy 1997;13:575-8.
8. Schaumkell JV, Villar RN. Healing of the ruptured ligamentum teres after hip dislocation—an arthroscopic finding. Hip Int 2009;19:64-6.
9. Wettstein M, Garofalo R, Borens O, Mouhsine E. Traumatic rupture of the ligamentum teres as a source of hip pain. Arthroscopy 2005;21:382.
10. Tschanner C, Hofmann S, Czerny C. [Hip dysplasia. Morphology, biomechanics and therapeutic principles with reference to the acetabular labrum]. Orthopade 1997;26: 89-108.
11. Altenberg AR. Acetabular labrum tears: a cause of hip pain and degenerative arthritis. South Med J 1977;70:174-5.
12. Cartledge JJ, Scott JH. The intumed acetabular labrum in osteoarthritis of the hip. J R Coll Surg Edinb 1982;27:339-44.
13. Crawford MJ, Dy CJ, Alexander JW, et al. The 2007 Frank Stinchfield Award. The bio-
14. Ferguson SJ, Bryant JT, Ganz R, Ito K. The influence of the acetabular labrum on hip joint cartilage consolidation: a poroelastic finite element model. J Biomech 2000;33:953-60.
15. Tschanner C, Hofmann S. [Labrum lesions in residual dysplasia of the hip joint. Biomechanical considerations on pathogenesis and treatment]. Orthopade 1998;27:725-32.
16. Petersen W, Petersen F, Tillmann B. Structure and vascularization of the acetabular labrum with regard to the pathogenesis and healing of labral lesions. Arch Orthop Trauma Surg 2003;123:283-8.
17. Putz R, Schrank C. [Anatomy of the labro-capsular complex]. Orthopade 1998;27:675-80.
18. McCarthy J, Noble P, Aluisio FV, et al. Anatomy, pathologic features, and treatment of acetabular labral tears. Clin Orthop Relat Res 2003;38-47.
19. Kelly BT, Shapiro GS, Digiovanni CW, et al. Vascularity of the hip labrum: a cadaveric investigation. Arthroscopy 2005;21:3-11.
20. Kim YT, Azuma H. The nerve endings of the acetabular labrum. Clin Orthop Relat Res 1995:176-81.
21. Ferguson SJ, Bryant JT, Ganz R, Ito K. An in vitro investigation of the acetabular labral seal in hip joint mechanics. J Biomech 2003;36:171-8.
22. Kachingwe AF, Grech S. Proposed algorithm for the management of athletes with athletic pubalgia (sports hernia): a case series. J Orthop Sports Phys Ther 2008;38:768-81.
23. Brannigan AE, Kerin MJ, McEntee GP. Gilmore's groin repair in athletes. J Orthop Sports Phys Ther 2000;30:329-32.
24. Bedi A, Dolan M, Leunig M, Kelly BT. Static and dynamic mechanical causes of hip pain. Arthroscopy 2011;27:235-51.
25. Farjo LA, Glick JM, Sampson TG. Hip arthroscopy for acetabular labral tears. Arthroscopy 1999;15:132-7.
26. Moorman CT 3rd, Warren RF, Hershman EB, et al. Traumatic posterior hip subluxation in American football. J Bone Joint Surg Am, 2003;85-A:1190-6.
27. Voos JE, Shindle MK, Pruett A, et al. Endoscopic repair of gluteus medius tendon tears of the hip. Am J Sports Med 2009;37:743-7.
28. Schilders E, Talbot JC, Robinson P, et al. Adductor-related groin pain in recreational athletes: role of the adductor enthesis, magnetic resonance imaging, and enthesal pubic cleft injections. J Bone Joint Surg Am 2009;91:2455-60.
29. Schilders E, Bismil Q, Robinson P, et al. Adductor-related groin pain in competitive athletes. Role of adductor enthesis, magnetic resonance imaging, and enthesal pubic cleft injections. J Bone Joint Surg Am 2007;89:2173-8.
30. Feeley BT, Powell JW, Muller MS, et al. Hip injuries and labral tears in the national

- football league. *Am J Sports Med* 2008;36:2187-95.
31. Markisz JA, Knowles RJ, Altchek DW, et al. Segmental patterns of avascular necrosis of the femoral heads: early detection with MR imaging. *Radiology* 1987;162:717-20.
 32. Mitchell DG, Rao VM, Dalinka MK, et al. Femoral head avascular necrosis: correlation of MR imaging, radiographic staging, radionuclide imaging, and clinical findings. *Radiology* 1987;162:709-15.
 33. Mont MA, Hungerford DS. Non-traumatic avascular necrosis of the femoral head. *J Bone Joint Surg Am* 1995;77:459-74.
 34. Schmitt-Sody M, Kirchhoff C, Mayer W, et al. Avascular necrosis of the femoral head: inter- and intraobserver variations of Ficat and ARCO classifications. *Int Orthop* 2008;32:283-7.
 35. Floerkemeier T, Thorey F, Daentzer D, et al. Clinical and radiological outcome of the treatment of osteonecrosis of the femoral head using the osteonecrosis intervention implant. *Int Orthop* 2011;35:489-95.
 36. Guo JJ, Tang N, Yang HL, et al. Impact of surgical approach on postoperative heterotopic ossification and avascular necrosis in femoral head fractures: a systematic review. *Int Orthop* 2010;34:319-22.
 37. Hsu HH, Rubin LJ. Iloprost inhalation solution for the treatment of pulmonary arterial hypertension. *Expert Opin Pharmacother* 2005;6:1921-30.
 38. M. Jäger, P. Hernigou, C. Zilkens, et al. [Cell therapy in bone-healing disorders]. *Orthopade* 2010;39:449-62.
 39. Jäger M, Tillmann FP, Thornhill TS, et al. Rationale for prostaglandin I₂ in bone marrow oedema—from theory to application. *Arthritis Res Ther* 2008;10:R120.
 40. Jäger M, Zilkens C, Bittersohl B, et al. Efficiency of iloprost treatment for osseous malperfusion. *Int Orthop*, 2011;35:761-5.
 41. Jäger M, Zilkens C, Westhoff B, et al. Efficiency of iloprost treatment for chemotherapy-associated osteonecrosis after childhood cancer. *Anticancer Res* 2009;29:3433-40.
 42. Ito K, Leunig M, Ganz R. Histopathologic features of the acetabular labrum in femoroacetabular impingement. *Clin Orthop Relat Res* 2004;262:71.
 43. Mintz DN, Hooper T, Connell D, et al. Magnetic resonance imaging of the hip: detection of labral and chondral abnormalities using noncontrast imaging. *Arthroscopy* 2005;21:385-93.
 44. Sundberg TP, Toomayan GA, Major NM. Toomayan, and N.M. Major, Evaluation of the acetabular labrum at 3.0-T MR imaging compared with 1.5-T MR arthrography: preliminary experience. *Radiology* 2006;238:706-11.
 45. Czerny C, Hofmann S, Neuhold A, et al. Lesions of the acetabular labrum: accuracy of MR imaging and MR arthrography in detection and staging. *Radiology* 1996;200:225-30.
 46. Locher S, Werlen S, Leunig M, Ganz R. [MR Arthrography with radial sequences for visualization of early hip pathology not visible on plain radiographs]. *Z Orthop Ihre Grenzgeb* 2002;140:52-7.
 47. Petersilge CA. MR arthrography for evaluation of the acetabular labrum. *Skeletal Radiol* 2001;30:423-30.
 48. Petersilge CA, Haque MA, Petersilge WJ, et al. Acetabular labral tears: evaluation with MR arthrography. *Radiology* 1996;200:231-5.
 49. Steinbach LS, Palmer WE, Schweitzer ME. Special focus session. MR arthrography. *Radiographics* 2002;22:1223-46.
 50. Keeney JA, Peelle MW, Jackson J, et al. Magnetic resonance arthrography versus arthroscopy in the evaluation of articular hip pathology. *Clin Orthop Relat Res* 2004;163-9.
 51. Czerny C, Oschatz E, Neuhold A, et al. [MR arthrography of the hip joint]. *Radiologe* 2002;42:451-6.
 52. Chan YS, Lien LC, Hsu HL, et al. Evaluating hip labral tears using magnetic resonance arthrography: a prospective study comparing hip arthroscopy and magnetic resonance arthrography diagnosis. *Arthroscopy* 2005;21:1250.
 53. Toomayan GA, Holman WR, Major NM, et al. Sensitivity of MR arthrography in the evaluation of acetabular labral tears. *AJR Am J Roentgenol* 2006;186:449-53.
 54. Leunig M, Werlen S, Ungersböck A, et al. Evaluation of the acetabular labrum by MR arthrography. *J Bone Joint Surg Br* 1997;79:230-4.
 55. Freedman BA, Potter BK, Dinanier PA, et al. Prognostic value of magnetic resonance arthrography for Czerny stage II and III acetabular labral tears. *Arthroscopy* 2006;22:742-7.
 56. Schmid MR, Nötzli HP, Zanetti M, et al. Cartilage lesions in the hip: diagnostic effectiveness of MR arthrography. *Radiology* 2003;226:382-6.
 57. Zlatkin MB, Pevsner D, Sanders TG, et al. Acetabular labral tears and cartilage lesions of the hip: indirect MR arthrographic correlation with arthroscopy—a preliminary study. *AJR Am J Roentgenol* 2010;194:709-14.
 58. Vahlensieck M, Peterfy CG, Wischer T, et al. Indirect MR arthrography: optimization and clinical applications. *Radiology* 1996;200:249-54.
 59. Kim YJ, Jaramillo D, Millis MB, et al. Assessment of early osteoarthritis in hip dysplasia with delayed gadolinium-enhanced magnetic resonance imaging of cartilage. *J Bone Joint Surg Am* 2003;85-A:1987-92.
 60. Jessel RH, Zilkens C, Tiderius C, et al. Assessment of osteoarthritis in hips with femoroacetabular impingement using delayed gadolinium enhanced MRI of cartilage. *J Magn Reson Imaging* 2009;30:1110-5.
 61. Jessel RH, Zurakowski D, Zilkens C, et al. Radiographic and patient factors associated with pre-radiographic osteoarthritis in hip dysplasia. *J Bone Joint Surg Am* 2009;91:1120-9.
 62. Pozzi G, Stradiotti P, Parra CG, et al. Femoro-acetabular impingement: can indirect MR arthrography be considered a valid method to detect endoarticular damage? A preliminary study. *Hip Int* 2009;19:386-91.
 63. Byrd JW, Jones KS. Diagnostic accuracy of clinical assessment, magnetic resonance imaging, magnetic resonance arthrography, and intra-articular injection in hip arthroscopy patients. *Am J Sports Med* 2004;32:1668-74.
 64. Nishii T, Nakanishi K, Sugano N, et al. Articular cartilage evaluation in osteoarthritis of the hip with MR imaging under continuous leg traction. *Magn Reson Imaging* 1998;16:871-5.
 65. Outerbridge RE. The etiology of chondromalacia patellae. *J Bone Joint Surg Br* 1961;43-B:752-7.
 66. Knuesel PR, Pfirrmann CW, Noetzi HP, et al. MR arthrography of the hip: diagnostic performance of a dedicated water-excitation 3D double-echo steady-state sequence to detect cartilage lesions. *AJR Am J Roentgenol* 2004;183:1729-35.
 67. Pfirrmann CW, Mengiardi B, Dora C, et al. Cam and pincer femoroacetabular impingement: characteristic MR arthrographic findings in 50 patients. *Radiology* 2006;240:778-85.
 68. Anderson LA, Peters CL, Park BB, et al. Acetabular cartilage delamination in femoroacetabular impingement. Risk factors and magnetic resonance imaging diagnosis. *J Bone Joint Surg Am* 2009;91:305-13.
 69. Beck M, Kalhor M, Leunig M, Ganz R. Hip morphology influences the pattern of damage to the acetabular cartilage: femoroacetabular impingement as a cause of early osteoarthritis of the hip. *J Bone Joint Surg Br* 2005;87:1012-8.
 70. Pfirrmann CW, Duc SR, Zanetti M, et al. MR arthrography of acetabular cartilage delamination in femoroacetabular cam impingement. *Radiology* 2008;249:236-41.
 71. Ganz R, Parvizi J, Beck M, et al. Femoroacetabular impingement: a cause for osteoarthritis of the hip. *Clin Orthop Relat Res* 2003;112-20.
 72. Reynolds D, Lucas J, Klaue K. Retroversion of the acetabulum. A cause of hip pain. *J Bone Joint Surg Br* 1999;81:281-8.
 73. Tonnies D, Heinecke A. Acetabular and femoral anteversion: relationship with osteoarthritis of the hip. *J Bone Joint Surg Am* 1999;81:1747-70.
 74. Siebenrock KA, Schoeniger R, Ganz R. Anterior femoro-acetabular impingement due to acetabular retroversion. Treatment with periacetabular osteotomy. *J Bone Joint Surg Am* 2003;85-A:278-86.
 75. Leunig M, Beck M, Dora C, Ganz R. [Femoroacetabular impingement: Trigger for the development of osteoarthritis.]. *Orthopade* 2006;35:77-84.
 76. Dora C, Zurbach J, Hersche O, Ganz R. Pathomorphologic characteristics of post-traumatic acetabular dysplasia. *J Orthop*

- Trauma 2000;14:483-9.
77. Dora C, Mascard E, Mladenov K, Seringe R. Retroversion of the acetabular dome after Salter and triple pelvic osteotomy for congenital dislocation of the hip. *J Pediatr Orthop B* 2002;11:34-40.
 78. Botser IB, Smith TW Jr, Nasser R, Domb BG. Open surgical dislocation versus arthroscopy for femoroacetabular impingement: a comparison of clinical outcomes. *Arthroscopy*, 2011;27:270-8.
 79. Dudda M, Albers C, Mamisch TC, et al. Do normal radiographs exclude asphericity of the femoral head-neck junction? *Clin Orthop Relat Res* 2009;467:651-9.
 80. Kubo T, Horii M, Harada Y, et al. Radial-sequence magnetic resonance imaging in evaluation of acetabular labrum. *J Orthop Sci* 1999;4:328-32.
 81. Ito K, Minka MA 2nd, Leunig M, et al. Femoroacetabular impingement and the cam-effect. A MRI-based quantitative anatomical study of the femoral head-neck offset. *J Bone Joint Surg Br* 2001;83:171-6.
 82. Siebenrock KA, Wahab KH, Werlen S, et al. Abnormal extension of the femoral head epiphysis as a cause of cam impingement. *Clin Orthop Relat Res* 2004;54:60.
 83. Beck M, Leunig M, Parvizi J, et al. Anterior femoroacetabular impingement: part II. Midterm results of surgical treatment. *Clin Orthop Relat Res* 2004;67-73.
 84. Kassarian A, Yoon LS, Belzile E, et al. Triad of MR arthrographic findings in patients with cam-type femoroacetabular impingement. *Radiology* 2005;236:588-92.
 85. Bittersohl B, Hosalkar HS, Apprich S, et al. Comparison of pre-operative dGEMRIC imaging with intra-operative findings in femoroacetabular impingement: preliminary findings. *Skeletal Radiol* 2011;40:553-61.
 86. Nötzli HP, Wyss TF, Stoecklin CH, et al. The contour of the femoral head-neck junction as a predictor for the risk of anterior impingement. *J Bone Joint Surg Br* 2002;84:556-60.
 87. Domayer SE, Mamisch TC, Kress I, et al. Radial dGEMRIC in developmental dysplasia of the hip and in femoroacetabular impingement: preliminary results. *Osteoarthritis Cartilage* 2010;18:1421-8.
 88. McCarthy JC, Lee JA. Acetabular dysplasia: a paradigm of arthroscopic examination of chondral injuries. *Clin Orthop Relat Res* 2002;122-8.
 89. Hodler J, Yu JS, Goodwin D, et al. MR arthrography of the hip: improved imaging of the acetabular labrum with histologic correlation in cadavers. *AJR Am J Roentgenol* 1995;165:887-91.
 90. Dinanuer PA, Murphy KP, Carroll JF. Sublabral sulcus at the posteroinferior acetabulum: a potential pitfall in MR arthrography diagnosis of acetabular labral tears. *AJR Am J Roentgenol* 2004;183:1745-53.
 91. Studler U, Kalberer F, Leunig M, et al. MR arthrography of the hip: differentiation between an anterior sublabral recess as a normal variant and a labral tear. *Radiology* 2008;249:947-54.
 92. Czerny C, Krestan C, Imhof H, et al. Magnetic resonance imaging of the postoperative hip. *Top Magn Reson Imaging* 1999;10:214-20.
 93. Chatha, D.S. and R. Arora, MR imaging of the normal hip. *Magn Reson Imaging Clin N Am* 2005;13:605-15.
 94. Klaue K, Durnin CW, Ganz R. The acetabular rim syndrome. A clinical presentation of dysplasia of the hip. *J Bone Joint Surg Br* 1991;73:423-9.
 95. DuBois DF, Omar IM. MR imaging of the hip: normal anatomic variants and imaging pitfalls. *Magn Reson Imaging Clin N Am* 2010;18:663-74.
 96. Fu Z, Peng M, Peng Q. Anatomical study of the synovial plicae of the hip joint. *Clin Anat* 1997;10:235-8.
 97. Blankenbaker DG, Davis KW, De Smet AA, Keene JS. MRI appearance of the pectinofoveal fold. *AJR Am J Roentgenol* 2009;192:93-5.
 98. Zilkens C, Jäger M, Bittersohl B, et al. [Slipped capital femoral epiphysis]. *Orthopade* 2010;39:1009-21.
 99. Zilkens J, Löer F, Zilkens KW. [Long-term Results After Conservatively Treated SCFE], in *Spätergebnisse in der Orthopädie*, Blauth/Ulrich, Editor. Springer: Berlin; 1986.
 100. Pitt MJ, Graham AR, Shipman JH, Birkby W. Herniation pit of the femoral neck. *AJR Am J Roentgenol* 1982;138:1115-21.
 101. Leunig M, Beck M, Kalhor M, et al. Fibrocystic changes at anterosuperior femoral neck: prevalence in hips with femoroacetabular impingement. *Radiology* 2005;236:237-46.
 102. Maroudas A, Evans H, Almeida L. Cartilage of the hip joint. Topographical variation of glycosaminoglycan content in normal and fibrillated tissue. *Ann Rheum Dis* 1973;32:1-9.
 103. Burstein D, Velyvis J, Scott KT, et al. Protocol issues for delayed Gd(DTPA)(2)-enhanced MRI (dGEMRIC) for clinical evaluation of articular cartilage. *Magn Reson Med* 2001;45:6-41.
 104. Tiderius CJ, Olsson LE, de Verdier H, et al. Gd-DTPA2)-enhanced MRI of femoral knee cartilage: a dose-response study in healthy volunteers. *Magn Reson Med* 2001;46:1067-71.
 105. Tiderius CJ, Olsson LE, Leander P, et al. Delayed gadolinium-enhanced MRI of cartilage (dGEMRIC) in early knee osteoarthritis. *Magn Reson Med* 2003;49:488-92.
 106. Watanabe A, Wada Y, Obata T, et al. Delayed gadolinium-enhanced MR to determine glycosaminoglycan concentration in reparative cartilage after autologous chondrocyte implantation: preliminary results. *Radiology* 2006;239:201-8.
 107. Bittersohl B, Hosalkar HS, Hughes T, et al. Feasibility of T2* mapping for the evaluation of hip joint cartilage at 1.5T using a three-dimensional (3D), gradient-echo (GRE) sequence: a prospective study. *Magn Reson Med* 2009;62:896-901.
 108. Trattnig S, Marlovits S, Gebetsroither S, et al. Three-dimensional delayed gadolinium-enhanced MRI of cartilage (dGEMRIC) for in vivo evaluation of reparative cartilage after matrix-associated autologous chondrocyte transplantation at 3.0T: Preliminary results. *J Magn Reson Imaging* 2007;26:974-82.
 109. Tiderius CJ, Jessel R, Kim YJ, Burstein D. Hip dGEMRIC in asymptomatic volunteers and patients with early osteoarthritis: the influence of timing after contrast injection. *Magn Reson Med* 2007;57:803-5.
 110. Cunningham T, Jessel R, Zurakowski D, et al. Delayed gadolinium-enhanced magnetic resonance imaging of cartilage to predict early failure of Bernese periacetabular osteotomy for hip dysplasia. *J Bone Joint Surg Am* 2006;88:1540-8.
 111. Miese FR, Zilkens C, Holstein A, et al. MRI morphometry, cartilage damage and impaired function in the follow-up after slipped capital femoral epiphysis. *Skeletal Radiol* 2010;39:533-41.
 112. Miese FR, Zilkens C, Holstein A, et al. Assessment of early cartilage degeneration after slipped capital femoral epiphysis using T2 and T2* mapping. *Acta Radiol* 2010.
 113. Zilkens C, Bittersohl B, Jäger M, et al. Significance of clinical and radiographic findings in young adults after slipped capital femoral epiphysis. *Int Orthop* 2010 [Epub ahead of print].
 114. Zilkens C, Holstein A, Bittersohl B, et al. Delayed gadolinium-enhanced magnetic resonance imaging of cartilage in the long-term follow-up after Perthes disease. *J Pediatr Orthop* 2010;30:147-53.
 115. Zilkens C, Miese F, Bittersohl B, et al. Delayed gadolinium-enhanced magnetic resonance imaging of cartilage (dGEMRIC), after slipped capital femoral epiphysis. *Eur J Radiol* 2010 [Epub ahead of print].
 116. Mamisch TC, Dudda M, Hughes T, et al. Comparison of delayed gadolinium enhanced MRI of cartilage (dGEMRIC) using inversion recovery and fast T1 mapping sequences. *Magn Reson Med* 2008;60:768-73.
 117. Bittersohl B, Hosalkar HS, Haamberg T, et al. Reproducibility of dGEMRIC in assessment of hip joint cartilage: a prospective study. *J Magn Reson Imaging* 2009;30:224-8.
 118. Bittersohl B, Hosalkar HS, Kim YJ, et al. T1 Assessment of Hip Joint Cartilage Following Intra-articular Gadolinium Injection: A Pilot Study. *Magn Reson Med* 2010;64:1200-7.
 119. Bittersohl B, Hosalkar HS, Werlen S, et al. Intravenous v/s Intraarticular Delayed Gadolinium-enhanced Magnetic Resonance Imaging (dGEMRIC) in the Hip Joint: A Comparative Analysis. *Invest Radiol* 2010;45:538-42.